PROVEN EFFICACY OF ACCELERATED DOUBLE-INJURY PROTOCOL: BIOPSY, ULTRASOUND, CLINICAL OBSERVATIONS

TWO-TREATMENT PROTOCOL FOR SKIN LAXITY USING 90-WATT DYNAMIC MONOPOLAR RADIOFREQUENCY DEVICE WITH REAL-TIME IMPEDANCE MONITORING


David McDaniel M.D.,¹ Robert Weiss M.D.,² Margaret Weiss M.D.,² Chris Mazur BS.¹ and Charrnaine Griffin CCR P¹

1. McDaniel Institute of Anti-Aging Research, Virginia Beach, VA; 2. Maryland Laser Skin & Vein Institute, Hunt Valley, MD

HIGHLIGHTS

- 24 female subjects enrolled and treated for facial skin laxity.
- 2 treatments with double treatment time instead of 4 standard treatments.
- Results show 35% reduction in skin laxity, 42% reduction in facial wrinkles and 33% reduction in the appearance of global photodamage.
- Histological analysis of biopsy samples proves significant increase in dermal collagen and elastin fibers.
- 19% increase in skin density evidenced by ultrasound measurements.

Pre



Post

Masson staining in one subject demonstrates increased Collagen (blue) at 3 months post treatment.